

GENERALIZED JACK POLYNOMIALS AND THE REPRESENTATION THEORY OF RATIONAL CHEREDNIK ALGEBRAS

CHARLES DUNKL AND STEPHEN GRIFFETH

ABSTRACT. We apply the Dunkl-Opdam operators and generalized Jack polynomials to study category $\mathcal{O}_c$ for the rational Cherednik algebra of type $G(r, p, n)$. We determine the set of aspherical values and, in case $p = 1$, answer a question of Iain Gordon on the ordering of category $\mathcal{O}_c$.

1. INTRODUCTION

We study the rational Cherednik algebra $\mathbb{H}_c$ of type $G(r, 1, n)$ by means of the Dunkl-Opdam operators introduced in [DuOp] and the generalized Jack polynomials introduced in [Gri]. Our main results are a characterization of the set of parameters c for which $\mathbb{H}_c e_\xi \mathbb{H}_c \neq \mathbb{H}_c$, where ξ is a linear character of $G(r, 1, n)$ and e_ξ is the corresponding idempotent, and a description, in terms of partition combinatorics, of an order on category $\mathcal{O}_c$ that arises by studying the generalized eigenspaces of the Dunkl-Opdam operators on standard modules.

Parameters c such that $\mathbb{H}_c e_\xi \mathbb{H}_c \neq \mathbb{H}_c$ are called ξ -aspherical, and it is a standard fact that the functor $M \mapsto e_\xi M$ gives an equivalence from $\mathbb{H}_c\text{-mod}$ to $e_\xi \mathbb{H}_c e_\xi\text{-mod}$ exactly if c is *not* ξ -aspherical. Writing e for the symmetrizing idempotent (corresponding to the trivial character), Proposition 5.5 of Berest-Chalykh [BeCh] states that for certain choices of ξ , we have an isomorphism $e \mathbb{H}_c e \cong e_\xi \mathbb{H}_{c'} e_\xi$ for a parameter c' depending on c and ξ , so combining this with our results allows one to establish Morita equivalences between rational Cherednik algebras at different parameters. We hope that by combining these equivalences with the techniques of the papers [Gor], [GoSt] and [GGS] one may obtain a tighter relationship between the Cherednik algebra of type $G(r, 1, n)$ and certain quiver varieties, including the Hilbert schemes of points on resolutions of type A surface singularities.

Our determination of the set of ξ -aspherical parameters is similar to that in [Dun] for the case $r = 1$; in this case the result was first proved by Gordon-Stafford [GoSt] and Bezrukavnikov-Etingof [BeEt]. The parameter c is a tuple $c = (c_0, d_0, \dots, d_{r-1})$ of complex numbers with $d_0 + d_1 + \dots + d_{r-1} = 0$ (see (1.4) for its relationship to $\mathbb{H}_c$). For any $l \in \mathbb{Z}$ we define $d_l = d_{l'}$ if $l = l' \bmod r$ and $0 \leq l' \leq r-1$. When $\xi = \text{triv}$ is the trivial character, we abbreviate ξ -aspherical values to *aspherical values*. Our first main result, Theorem 3.4, implies that the set of aspherical values is a certain finite union of hyperplanes. Here we restate it with the minimum amount of notation.

Theorem 1.1. *The parameter c is aspherical exactly if*

- (a) $c_0 = -k/m$ for integers k and m satisfying $1 \leq k < m \leq n$ or
- (b) there is an integer $0 \leq l \leq r-1$, an integer $-(n-1) \leq m \leq n-1$, and an integer k such that $k \not\equiv 0 \bmod r$,

$$k = d_l - d_{l-k} + r m c_0, \quad \text{and} \quad 1 \leq k \leq l + \left(\sqrt{n + \frac{1}{4}m^2} - \frac{1}{2}m - 1 \right) r$$

We are very grateful to Pavel Etingof for pointing out Lemma 3.3, which helped to put us on the right track, to an anonymous referee for careful reading and helpful advice, and to Iain Gordon for stimulating conversation and useful input. The second author acknowledges the full financial support of EPSRC grant EP/G007632.

As in Lemma 3.3, the hyperplanes of part (b) may also be described as follows: for each integer $0 \leq l \leq r-1$, each rectangular partition λ with at most n boxes, and each integer $k \not\equiv 0 \pmod r$ with

$$1 \leq k \leq l + r(\text{length}(\lambda) - 1)$$

the hyperplane

$$k = d_l - d_{l-k} + r(\lambda_1 - \text{length}(\lambda))c_0$$

is aspherical. See Corollary 3.5 for other linear characters ξ of $G(r, 1, n)$, and Corollary 3.6 for the case of $G(r, p, n)$.

The irreducible representations of $G(r, 1, n)$ are in sensible bijection with r -partitions $\lambda^\bullet = (\lambda^0, \dots, \lambda^{r-1})$ of n , and attached to each irreducible representation $S^{\lambda^\bullet}$ of $G(r, 1, n)$ is a *standard module* $M_c(\lambda^\bullet)$ of $\mathbb{H}_c$. The proof of Theorem 3.4 consists of computing the norm, with respect to the contravariant form, of a certain invariant polynomial in each standard module. This gives a necessary condition for c to be aspherical. On the other hand we can show that all such parameters are aspherical using Theorem 7.5 of [Gri] and the results of [BeEt].

By using the calculation of the spectrum of standard modules for $\mathbb{H}_c$ in Theorem 5.1 of [Gri], we also answer question 10.1 from [Gor]. Gordon's question has to do with *category* $\mathcal{O}_c$ for the Cherednik algebra: this is the Serre subcategory of $\mathbb{H}_c$ -modules generated by the standard modules $M_c(\lambda^\bullet)$. Category $\mathcal{O}_c$ is a highest weight category with standard objects the modules $M_c(\lambda^\bullet)$, where the order comes from a certain grading element of $\mathbb{H}_c$ (the *deformed Euler field*).

When $d_l/rc_0 \in \mathbb{Z}$ for $0 \leq l \leq r-1$ the tuple

$$(d_{r-1}/rc_0, \dots, d_0/rc_0)$$

is an element of the root lattice of type A_{r-1} and hence corresponds, according to the (standard) bijection described in 4.2, to an r -core partition μ_c . Via the bijection between ordinary partitions with r -core μ_c and r -partitions obtained by taking r -quotients, dominance order on ordinary partitions induces an ordering $\geq'_c$ on r -partitions. This ordering arose geometrically from quiver varieties in [Gor]; here we show that it arises also from the spectrum of the Dunkl-Opdam subalgebra. It should be quite interesting to investigate a conceptual reason for this coincidence.

For any parameter $c = (c_0, d_0, \dots, d_{r-1}) \in \mathbb{R}^{r+1}$ with $c_0 > 0$, we define, as in (4.1), another ordering on r -partitions $\lambda^\bullet = (\lambda^0, \dots, \lambda^{r-1})$ by $\lambda^\bullet \geq_c \chi^\bullet$ if for all $j \in \mathbb{R}$ and $0 \leq l \leq r-1$,

$$\begin{aligned} & |\{b \in \lambda^\bullet \mid \frac{d_{\beta(b)}}{rc_0} + \text{ct}(b) > j \text{ or } \frac{d_{\beta(b)}}{rc_0} + \text{ct}(b) = j \text{ and } \beta(b) \leq l\}| \\ & \geq |\{b' \in \chi^\bullet \mid \frac{d_{\beta(b')}}{rc_0} + \text{ct}(b') > j \text{ or } \frac{d_{\beta(b')}}{rc_0} + \text{ct}(b') = j \text{ and } \beta(b') \leq l\}|. \end{aligned}$$

For the definitions of the statistics $\text{ct}(b)$ and $\beta(b)$ see (1.9) and (1.10). With these notations, Theorem 4.1 and Corollary 4.3 state that

Theorem 1.2. (a) *For any choice of parameter $c \in \mathbb{R}^{r+1}$ with $c_0 > 0$, category $\mathcal{O}_c$ is a highest weight category with respect to the ordering $\geq_c$ described above;*
 (b) *if $d_l/rc_0 \in \mathbb{Z}$ for $0 \leq l \leq r-1$ then $\mathcal{O}_c$ is a highest weight category with respect to the ordering $\geq'_c$ described above.*

Since there are fewer order relations for these orders than for the order coming from the deformed Euler field, our result broadens the applicability of Theorem 4.49 of [Rou], allowing one to deduce Morita equivalences between category $\mathcal{O}$'s at different parameters.

1.1. The symmetric group. Let S_n be the group of permutations of the set $\{1, 2, \dots, n\}$. The notation $w_1 \leq w_2$ for $w_1, w_2 \in S_n$ refers to Bruhat order, and we write $l(w)$ for the length of an element $w \in S_n$. Let $w_0 \in S_n$ be the *longest element*, with

$$(1.1) \quad w_0(i) = n - i + 1 \quad \text{for } 1 \leq i \leq n.$$

For a sequence $\mu \in \mathbb{Z}_{\geq 0}^n$ of n non-negative integers, we write μ^+ for the non-increasing (partition) rearrangement of μ , and μ^- for the non-decreasing (anti-partition) rearrangement of μ . For $w \in S_n$ and $\mu \in \mathbb{Z}_{\geq 0}^n$, the formula

$$(1.2) \quad w \cdot \mu = (\mu_{w^{-1}(1)}, \mu_{w^{-1}(2)}, \dots, \mu_{w^{-1}(n)})$$

defines a left action of S_n on $\mathbb{Z}_{\geq 0}^n$. Let w_μ be the longest element of S_n such that $w_\mu \cdot \mu = \mu^-$; thus

$$(1.3) \quad w_\mu(i) = |\{1 \leq j < i \mid \mu_j < \mu_i\}| + |\{i \leq j \leq n \mid \mu_j \leq \mu_i\}|.$$

Also define the *rank function* r_μ by

$$(1.4) \quad r_\mu(i) = |\{1 \leq j \leq i \mid \mu_j \geq \mu_i\}| + |\{i < j \leq n \mid \mu_j > \mu_i\}|$$

so that

$$(1.5) \quad w_\mu(i) + r_\mu(i) = n + 1 \quad \text{or equivalently} \quad w_\mu = w_0 r_\mu.$$

There is a partial order $\geq$ on $\mathbb{Z}_{\geq 0}^n$ defined as follows: for $\mu, \nu \in \mathbb{Z}_{\geq 0}^n$ we have

$$(1.6) \quad \mu > \nu \quad \text{if} \quad \mu_+ >_d \nu_+ \quad \text{or} \quad \mu_+ = \nu_+ \quad \text{and} \quad w_\mu > w_\nu,$$

where $>_d$ is dominance order on partitions. For a reminder of the definition of dominance order on partitions, see (4.11); here we use the notation $>_d$ to distinguish it from the order on $\mathbb{Z}_{\geq 0}^n$ that we are defining.

1.2. Partitions, Young diagrams, and tableaux. Let $n \in \mathbb{Z}_{>0}$ be a positive integer. A *partition of length n* is a non-increasing sequence $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n > 0)$ of n positive integers. Without reference to length, a *partition* is a non-increasing sequence $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_m \geq 0)$ of non-negative integers, and we consider two partitions to be equal if they differ by a terminal string of zeros. In particular, we refer to a sequence of 0's as the *empty partition* $\emptyset$. Let r be a positive integer. An *r -partition* is a sequence $\lambda^\bullet = (\lambda^0, \lambda^1, \dots, \lambda^{r-1})$ (some λ^i 's may be empty) of r partitions. The *Young diagram* of an r -partition is the graphical representation consisting of a collection of boxes stacked in a corner: the Young diagram for the 4-partition $((3, 3, 1), (2, 1), \emptyset, (5, 5, 2, 1))$ of 23 is

$$\left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \\ \hline \end{array}, \emptyset, \begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \square & \square & \square & \square & \square \\ \hline \square & \square & & & \\ \hline \square & & & & \\ \hline \end{array} \right).$$

A *tableau* T on an r -partition $\lambda^\bullet$ of n is a filling of the boxes of $\lambda^\bullet$ with non-negative integers. A tableau T is *column-strict* if within each partition λ^i , its entries are strictly increasing from top to bottom and weakly increasing from left to right. A *standard Young tableau* on $\lambda^\bullet$ is a bijection T from the boxes of $\lambda^\bullet$ to the set $\{1, 2, \dots, n\}$ in such a way that the entries are increasing left to right and top to bottom. For example,

$$\left(\begin{array}{|c|c|c|} \hline 2 & 4 & 6 \\ \hline 3 & 9 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 5 \\ \hline 7 & 8 \\ \hline \end{array} \right)$$

is a standard Young tableau on the 2-partition $((3, 2), (2, 2))$. We write

$$(1.7) \quad T(b) = i \quad \text{if } i \text{ appears in box } b.$$

Then for standard Young tableaux S and T on λ , the permutation TS^{-1} of $\{1, 2, \dots, n\}$ is a measure of the distance between S and T . For an r -partition $\lambda^\bullet$ we write

$$(1.8) \quad \text{SYT}(\lambda) = \{\text{standard Young tableaux on } \lambda^\bullet\}$$

For a box $b \in \lambda$, write $\text{row}(b) = i$ and $\text{col}(b) = j$ if b is in the i th row and j th column of λ , and define the *content* $\text{ct}(b)$ of b by

$$(1.9) \quad \text{ct}(b) = \text{col}(b) - \text{row}(b).$$

We also define the function β on the set of boxes of $\lambda^\bullet$ by

$$(1.10) \quad \beta(b) = l \quad \text{if } b \in \lambda^l.$$

Thus for the tableau T pictured above, one has

$$\beta(T^{-1}(7)) = 1 \quad \text{and} \quad \text{ct}(T^{-1}(3)) = -1.$$

1.3. The groups $G(r, 1, n)$. Fix positive integers r and n . A *monomial matrix* is a square matrix with exactly one non-zero entry in each row and each column. Let

$$(1.11) \quad W = G(r, 1, n)$$

be the group of n by n monomial matrices whose non-zero entries are r th roots of 1. We write $\mathbb{C}W$ for the complex group algebra of W . Let $\zeta = e^{2\pi i/r}$ and

$$(1.12) \quad \zeta_i = \text{diag}(1, \dots, \zeta, \dots, 1)$$

be the diagonal matrix with a ζ in the i th position and 1's elsewhere on the diagonal. Let

$$(1.13) \quad s_{ij} = (ij) \quad \text{and} \quad s_i = s_{i,i+1}$$

be the transposition interchanging i and j and the i th simple transposition, respectively. We write $\mathfrak{h} = \mathbb{C}^n$ for the defining representation of W and $\mathfrak{h}^*$ for its dual.

There is a version of Young's orthonormal form that works for the groups $G(r, 1, n)$: the irreducible complex representations of $G(r, 1, n)$ may be indexed by the r -partitions of n in such a way that if $S^{\lambda^\bullet}$ is the irreducible representation corresponding to the r -partition $\lambda^\bullet$, then $S^{\lambda^\bullet}$ has a basis v_T indexed by the set of standard Young tableaux T on $\lambda^\bullet$. The vectors v_T are eigenvectors for a certain maximal commutative subalgebra of $\mathbb{C}W$, and the action of the generators ζ_i and s_i on v_T can be made quite explicit: see section 3 of [Gri] for these facts.

1.4. The rational Cherednik algebra. Let c_0 and d_i for $i \in \mathbb{Z}$ be variables such that $d_i = d_j$ if $i = j \bmod r$, and $d_0 + d_1 + \dots + d_{r-1} = 0$. Let $k = \mathbb{C}[c_0, d_i]_{1 \leq i \leq r-1}$ be the polynomial ring over $\mathbb{C}$ generated by these variables and let F be its fraction field. The *rational Cherednik algebra* $\mathbb{H}$ for $W = G(r, 1, n)$ is generated by $k[x_1, \dots, x_n]$, kW , and $k[y_1, \dots, y_n]$ with relations

$$(1.14) \quad y_i x_i = x_i y_i + 1 - c_0 \sum_{j \neq i} \sum_{l=0}^{r-1} \zeta_i^l s_{ij} \zeta_i^{-l} - \sum_{j=0}^{r-1} (d_j - d_{j-1}) e_{ij}$$

and for $i \neq j$

$$(1.15) \quad y_i x_j = x_j y_i + c_0 \sum_{l=0}^{r-1} \zeta^{-l} \zeta_i^l s_{ij} \zeta_i^{-l},$$

where for $1 \leq i \leq n$ and $0 \leq j \leq r-1$

$$(1.16) \quad e_{ij} = \frac{1}{r} \sum_{l=0}^{r-1} \zeta^{-lj} \zeta_i^l$$

are the idempotents for the cyclic reflection subgroups of W . These parameters are related to those found in Gordon's paper [Gor] by the equations

$$(1.17) \quad H_j = \frac{1}{r} (d_{j-1} - d_j) \quad \text{for } 0 \leq j \leq r-1, \text{ and } h = -c_0,$$

and to those in section 6 of Rouquier's paper [Rou] by

$$(1.18) \quad h_j = -\frac{1}{r}d_j \quad \text{for } 0 \leq j \leq r-1, \text{ and } h = -c_0.$$

Thus setting $q = e^{-2\pi i c_0}$ and $Q_j = e^{-2\pi i d_j/r}$ for $0 \leq j \leq r-1$ one obtains the parameters for the corresponding finite Hecke algebra of type $G(r, 1, n)$. Assuming $r > 1$ and $n > 1$ as in 6.1.2 of [Rou], this finite Hecke algebra has generators $T_0, T_1, \dots, T_{n-1}$, with the relations $T_0 T_1 T_0 T_1 = T_1 T_0 T_1 T_0$, braid relations between T_i and T_j for $1 \leq i < j \leq n-1$, and

$$(T_i - q)(T_i + 1) = 0 \quad \text{for } 1 \leq i \leq n-1, \text{ and } \prod_{j=0}^{r-1} (T_0 - e^{2\pi i j/r} Q_j) = 0.$$

The *PBW theorem* for $\mathbb{H}$ asserts that as a k -module

$$(1.19) \quad \mathbb{H} \cong k[x_1, \dots, x_n] \otimes kW \otimes k[y_1, \dots, y_n].$$

It implies that if we define the *standard module* $M(\lambda^\bullet)$ by

$$(1.20) \quad M(\lambda^\bullet) = \text{Ind}_{k[y_1, \dots, y_n] \rtimes kW}^{\mathbb{H}} S^{\lambda^\bullet}$$

then as a $k[x_1, \dots, x_n] \rtimes kW$ -module,

$$(1.21) \quad M(\lambda^\bullet) \cong k[x_1, \dots, x_n] \otimes S^{\lambda^\bullet}.$$

Let $\bar{\cdot}$ be the automorphism of k that fixes $c_0, d_0, \dots, d_{r-1}$ and acts as complex conjugation on $\mathbb{C}$. Let $x \mapsto \bar{x}$ be a conjugate-linear W -equivariant isomorphism from $\mathfrak{h}^*$ onto $\mathfrak{h}$. Fix a W -invariant positive definite Hermitian form on $S^{\lambda^\bullet}$. Then $M(\lambda^\bullet)$ carries a *contravariant form* $\langle \cdot, \cdot \rangle$, determined uniquely by the requirements that it be linear in the second variable, skew-symmetric, and satisfy the conditions

$$(1.22) \quad \langle w.f, w.g \rangle = \langle f, g \rangle \quad \text{and} \quad \langle x.f, g \rangle = \langle f, \bar{x}.g \rangle \quad \text{for } f, g \in M(\lambda^\bullet), w \in W, \text{ and } x \in \mathfrak{h}^*,$$

and restrict to the given form on $S^{\lambda^\bullet}$. The proof of Theorem 2.18 of [DuOp] works, *mutatis mutandis*, for this situation.

If we choose a particular specialization $c_0, d_i \in \mathbb{C}$ of the parameters to complex numbers, we write $\mathbb{H}_c$ for the resulting algebra. The specializations $M_c(\lambda^\bullet)$ of the standard modules have unique irreducible quotients $L_c(\lambda^\bullet)$, and we define *category* $\mathcal{O}_c$ to be the category of finitely generated $\mathbb{H}_c$ -modules on which $y_1, \dots, y_n$ act locally nilpotently. The contravariant form also specializes to a conjugate-symmetric form $\langle \cdot, \cdot \rangle_c$, provided that the parameters are real numbers. Otherwise it is not conjugate-symmetric; see Theorem 2.18 of [DuOp].

1.5. The Dunkl-Opdam subalgebra and the non-symmetric generalized Jack polynomials. For $1 \leq i \leq n$ define

$$(1.23) \quad z_i = y_i x_i + c_0 \phi_i \quad \text{where } \phi_i = \sum_{1 \leq j < i} \zeta_i^l s_{ij} \zeta_i^{-l} \text{ are the Jucys-Murphy elements for } G(r, 1, n).$$

The elements z_i were introduced in [DuOp], where it is also proved that they commute. The *Dunkl-Opdam* subalgebra $\mathfrak{t}$ of $\mathbb{H}$ is generated by $z_1, \dots, z_n$ and $\zeta_1, \dots, \zeta_n$. By the PBW theorem it is isomorphic to the polynomial ring in the variables $z_1, \dots, z_n$ tensored with the group algebra of $(\mathbb{Z}/r\mathbb{Z})^n$. Since the operators z_i are fixed by the conjugate-linear anti-involution $\bar{\cdot}$ of $\mathbb{H}$ such that $\bar{x}_i = y_i$ and $\bar{w} = w^{-1}$ for all $1 \leq i \leq n$ and $w \in G(r, 1, n)$, they act as self-adjoint operators on $M(\lambda^\bullet)$ with respect to the contravariant form.

The following theorem combines Theorems 5.1 and 6.1 of [Gri]. The T^{-1} 's appear here because we are using the "inverse" definition of standard Young tableau to that in [Gri]. Define a partial order on pairs $(\mu, T) \in \mathbb{Z}_{\geq 0}^n \times \text{SYT}(\lambda^\bullet)$ by $(\mu, T) > (\nu, S)$ exactly if $\mu > \nu$, where the order on $\mathbb{Z}_{\geq 0}^n$ is defined by (1.6).

Theorem 1.3. Let $\lambda^\bullet$ be an r -partition of n , $\mu \in \mathbb{Z}_{\geq 0}^n$, and let T be a standard tableau on $\lambda^\bullet$. Put $v_T^\mu = w_\mu^{-1}.v_T$ and recall the definitions of β and ct given in (1.9) and (1.10).

(a) The action of ζ_i and z_i on $M(\lambda^\bullet)$ are given by

$$\zeta_i.x^\mu v_T^\mu = \zeta^{\beta(T^{-1}w_\mu(i))-\mu_i} x^\mu v_T^\mu$$

and

$$\begin{aligned} z_i.x^\mu v_T^\mu &= (\mu_i + 1 - (d_{\beta(T^{-1}w_\mu(i))} - d_{\beta(T^{-1}w_\mu(i))-\mu_i-1}) - r \text{ct}(T^{-1}w_\mu(i))c_0) x^\mu v_T^\mu \\ &\quad + \sum_{(\nu, S) < (\mu, T)} c_{\nu, S} x^\nu v_S^\nu. \end{aligned}$$

(b) Assuming that scalars are extended to $F = \mathbb{C}(c_0, d_1, d_2, \dots, d_{r-1})$, for each $\mu \in \mathbb{Z}_{\geq 0}^n$ and $T \in \text{SYT}(\lambda^\bullet)$ there exists a unique $\mathfrak{t}$ eigenvector $f_{\mu, T} \in M(\lambda)$ such that

$$f_{\mu, T} = x^\mu v_T^\mu + \text{lower terms.}$$

The $\mathfrak{t}$ -eigenvalue of $f_{\mu, T}$ is determined by the formulas in part (a).

(c) Write

$$a_i = \text{ct}(T^{-1}w_\mu(i)) \quad \text{and} \quad b_i = \beta(T^{-1}w_\mu(i)).$$

Then the norm of $f_{\mu, T}$ is given by

$$\begin{aligned} \langle f_{\mu, T}, f_{\mu, T} \rangle &= \prod_{i=1}^n \prod_{k=1}^{\mu_i} (k - (d_{b_i} - d_{b_i-k}) - r a_i c_0) \\ &\quad \times \prod_{\substack{1 \leq i < j \leq n \\ \mu_i > \mu_j}} \prod_{\substack{1 \leq k \leq \mu_i - \mu_j \\ k = b_i - b_j \pmod r}} \frac{(k - (d_{b_i} - d_{b_j}) - r(a_i - a_j)c_0)^2 - (r c_0)^2}{(k - (d_{b_i} - d_{b_j}) - r(a_i - a_j)c_0)^2} \\ &\quad \times \prod_{\substack{1 \leq i < j \leq n \\ \mu_i < \mu_j - 1}} \prod_{\substack{1 \leq k \leq \mu_j - \mu_i - 1 \\ k = b_j - b_i \pmod r}} \frac{(k - (d_{b_j} - d_{b_i}) - r(a_j - a_i)c_0)^2 - (r c_0)^2}{(k - (d_{b_j} - d_{b_i}) - r(a_j - a_i)c_0)^2}. \end{aligned}$$

2. GENERALIZED JACK POLYNOMIALS

2.1. Definition of the (symmetric) generalized Jack polynomials. The functions $f_{\mu, T}$ are a basis of $M(\lambda^\bullet)$. Let $e = \sum_{w \in S_n} w$ be the S_n -symmetrizer. Then the functions $e.f_{\mu, T}$ such that $\mu_i = \beta(T^{-1}w_\mu(i)) \pmod r$ for $1 \leq i \leq n$ span the $G(r, 1, n)$ invariants in $M(\lambda^\bullet)$. To single a basis out from this spanning set we use the following construction from [Dun].

Given $(\mu, T) \in \mathbb{Z}_{\geq 0}^n \times \text{SYT}(\lambda^\bullet)$, define a function $S = S(\mu, T)$ on the boxes of $\lambda^\bullet$ by $S(b) = \mu_{w_\mu^{-1}T(b)}$. It follows from the definition that S is weakly increasing top to bottom and left to right. For instance, if

$$\mu = (2, 3, 2, 0, 4, 2, 5, 2, 2) \quad \text{with} \quad w_\mu = (6, 7, 5, 1, 8, 4, 9, 3, 2)$$

and

$$T = \left(\begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 8 & 9 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & 6 \\ \hline 5 & 7 \\ \hline \end{array} \right) \quad \text{then} \quad S(\mu, T) = \left(\begin{array}{|c|c|c|} \hline 0 & 2 & 2 \\ \hline 4 & 5 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 2 & 3 \\ \hline \end{array} \right).$$

One checks that $\mu_i = \beta(T^{-1}w_\mu(i)) \pmod r$ for all $1 \leq i \leq n$ exactly if $\beta(b) = S(b) \pmod r$ for all $b \in \lambda^\bullet$. Therefore the invariants in $M(\lambda^\bullet)$ for the diagonal subgroup of $G(r, 1, n)$ generated by $\zeta_1, \dots, \zeta_n$ are those $f_{\mu, T}$ so that $S(b) = \beta(b) \pmod r$ for all $b \in \lambda^\bullet$, where $S = S(\mu, T)$. We will describe the $G(r, 1, n)$ invariants.

The (Cherednik-style) intertwining operators are

$$(2.1) \quad \sigma_i = s_i + f_{\alpha_i}, \quad \text{where} \quad f_{\alpha_i} = \frac{c_0}{z_i - z_{i+1}} \sum_{l=0}^{r-1} \zeta_i^l \zeta_{i+1}^{-l}.$$

Direct calculation shows the σ_i 's satisfy the braid relations, and that

$$(2.2) \quad \sigma_i^2 = 1 - f_{\alpha_i}^2$$

as in Lemma 5.2 (a) of [Gri2]. For a permutation $w \in S_n$ written as a reduced word in the simple reflections $w = s_{i_1} \cdots s_{i_p}$ we define $\sigma_w = \sigma_{i_1} \cdots \sigma_{i_p}$; by the braid relations this does not depend on our choice of reduced word for w . It follows from their definition that the intertwiners are self-adjoint for the contravariant form on $M(\lambda^\bullet)$.

- Lemma 2.1.** (a) If S is a column-strict tableau on $\lambda^\bullet$ then there is a non-decreasing $\mu \in \mathbb{Z}_{\geq 0}^n$ and $T \in \text{SYT}(\lambda^\bullet)$ with $S = S(\mu, T)$.
 (b) The F -span of the polynomials $e.f_{\mu, T}$ as T ranges over $\text{SYT}(\lambda^\bullet)$ and μ ranges over non-decreasing sequences is all of $M(\lambda^\bullet)^{S_n}$.
 (c) If $\mu \in \mathbb{Z}_{\geq 0}^n$ is non-decreasing and $T_1, T_2 \in \text{SYT}(\lambda^\bullet)$ with $S(\mu, T_1) = S(\mu, T_2)$ then $e.f_{\mu, T_1}$ and $e.f_{\mu, T_2}$ are $\mathbb{C}$ -multiples of one another.
 (d) If $\mu \in \mathbb{Z}_{\geq 0}^n$ is non-decreasing, $T \in \text{SYT}(\lambda^\bullet)$ and $S(\mu, T)$ is not column-strict then $e.f_{\mu, T} = 0$.

Proof. Part (a) follows from the definitions given above. For part (b), the formula given in part (a) of Lemma 5.3 of [Gri] implies that for each $(\nu, T) \in \mathbb{Z}_{\geq 0}^n \times \text{SYT}(\lambda^\bullet)$ we have $f_{\nu, T} = \sigma_w f_{\nu^-, T}$ for some $w \in S_n$. But $\sigma_w.f_{\nu^-, T} \in FG(r, 1, n)f_{\nu^-, T}$ by definition of σ_w , so $e.f_{\nu, T}$ is an F -multiple of $e.f_{\nu^-, T}$, proving (b).

For (c), assuming $S(\mu, T_1) = S(\mu, T_2)$ we obtain $R(\mu, T_1) = R(\mu, T_2)$ where $R(\mu, T)$ is defined in 7.17 of [Gri]. Hence Lemma 7.4 of [Gri] shows that there is a sequence of intertwiners $\sigma_{i_1}, \dots, \sigma_{i_l}$ with $f_{\mu, T_1} = \sigma_{i_l} \cdots \sigma_{i_1} f_{\mu, T_2}$ and so that $\sigma_{i_j} \cdots \sigma_{i_1} f_{\mu, T_2} = f_{\mu, S_j}$ for standard Young tableaux S_j with the property that $R(\mu, S_j) = R(\mu, T_2)$ for all $1 \leq j \leq l$. We claim that $f_{\mu, S_j} \in \mathbb{C}S_n.f_{\mu, T_2}$ for all j ; part (c) follows immediately from this. We may therefore assume $l = 1$, so that $f_{\mu, T_1} = \sigma_i.f_{\mu, T_2}$ and $\mu_i = \mu_{i+1}$. But $\sigma_i = s_i + f_{\alpha_i}$, and the condition $\mu_i = \mu_{i+1}$ together with the formula in Theorem 1.3 for the $\mathfrak{t}$ -eigenvalue of f_{μ, T_2} imply that f_{α_i} acts by a complex number on f_{μ, T_2} , finishing the proof of (c).

We now prove (d). Assuming $S = S(\mu, T)$ is not column-strict, there are boxes $b, b' \in \lambda^\bullet$ so that b' is directly below b and $S(b') = S(b)$. By part (c) we may prove (d) for any tableau T' with $S(\mu, T) = S(\mu, T')$. We may therefore assume that $w_\mu^{-1}T(b) = i + 1$ and $w_\mu^{-1}T(b') = i$ for some integer $1 \leq i \leq n - 1$. Using part (c) of Lemma 5.3 of [Gri] implies $\sigma_i.f_{\mu, T} = 0$, and the formula in Theorem 1.3 for the $\mathfrak{t}$ -weight of $f_{\mu, T}$ implies that $s_i.f_{\mu, T} = -f_{\mu, T}$ whence $e.f_{\mu, T} = es_i.f_{\mu, T} = -e.f_{\mu, T}$. Thus $e.f_{\mu, T} = 0$. $\square$

For each column-strict tableau S on $\lambda^\bullet$ with $S(b) = \beta(b) \bmod r$ let $\mu \in \mathbb{Z}_{\geq 0}^n$ be the sequence obtained by arranging the entries of S in non-decreasing order, and fix a tableau $T \in \text{SYT}(\lambda^\bullet)$ with $S = S(\mu, T)$. Define the *generalized Jack polynomial* by

$$(2.3) \quad g_S = e.f_{\mu, T}.$$

The definition of g_S depends on the choice of T , but only up to multiplication by a constant in $\mathbb{C}$.

2.2. The norm of g_S . In the following theorem, and all that follows, we use the convention that the product over an empty set is 1.

Theorem 2.2. *The functions g_S , where S ranges over column strict tableau satisfying $S(b) = \beta(b) \bmod r$ for all $b \in \lambda^\bullet$, are a basis of $M(\lambda^\bullet)^W$. The norm of g_S is given by*

$$\begin{aligned} \langle g_S, g_S \rangle &= n! \prod_{b \in \lambda^\bullet} \prod_{1 \leq k \leq S(b)} (k - (d_{\beta(b)} - d_{\beta(b)-k}) - r \text{ct}(b) c_0) \\ &\quad \times \prod_{b, b' \in \lambda^\bullet} \prod_{\substack{1 \leq k \leq S(b) - S(b') \\ k = \beta(b) - \beta(b') \bmod r}} \frac{k - (d_{\beta(b)} - d_{\beta(b')}) - r(\text{ct}(b) - \text{ct}(b') - 1) c_0}{k - (d_{\beta(b)} - d_{\beta(b')}) - r(\text{ct}(b) - \text{ct}(b')) c_0} \\ &\quad \times \prod_{b, b' \in \lambda^\bullet} \prod_{\substack{1 \leq k \leq S(b) - S(b') - r \\ k = \beta(b) - \beta(b') \bmod r}} \frac{k - (d_{\beta(b)} - d_{\beta(b')}) - r(\text{ct}(b) - \text{ct}(b') + 1) c_0}{k - (d_{\beta(b)} - d_{\beta(b')}) - r(\text{ct}(b) - \text{ct}(b')) c_0} \end{aligned}$$

Proof. We first show that the functions g_S have distinct $\mathfrak{t}^{S_n}$ eigenvalues; the fact that they are linearly independent follows from this. The eigenvalues of symmetric functions in $z_1, \dots, z_n$ acting on g_S are determined by the multiset of eigenvalues of $z_1, \dots, z_n$ acting on any $f_{\mu, T}$ for which $S = S(\mu, T)$. By the definition of $S(\mu, T)$ given above and part (a) of Theorem 1.3 this multiset is $\{\mu_i + 1 - (d_{\beta(T^{-1}w_\mu(i))} - d_{-1}) - r \text{ct}(T^{-1}w_\mu(i)) c_0 \mid 1 \leq i \leq n\} = \{S(b) + 1 - (d_{\beta(b)} - d_{-1}) - r \text{ct}(b) c_0 \mid b \in \lambda^\bullet\}$ and S is uniquely determined by it, because any filling of a partition which is weakly increasing left to right and top to bottom is uniquely determined by the associated multiset of contents.

We next calculate the norm of $g_S = e.f_{\mu, T}$ using a version of the argument proving (5.8.24) in [Mac2]. We write $R = \{\epsilon_i - \epsilon_j \mid 1 \leq i \neq j \leq n\}$ for the root system of type A_{n-1} and $R^+ = \{\epsilon_i - \epsilon_j \mid 1 \leq i < j \leq n\}$ for the set of positive roots. For a root $\alpha = \epsilon_i - \epsilon_j$ of S_n write

$$(2.4) \quad f_\alpha = \frac{c_0}{z_i - z_j} \sum_{l=0}^{r-1} (\zeta_i \zeta_j^{-1})^l$$

and let g_α be the constant by which f_α acts on $f_{\mu, T}$. This is well-defined by the column-strictness of S : setting $b_1 = T^{-1}w_\mu(i)$ and $b_2 = T^{-1}w_\mu(j)$, f_α acts by 0 on $f_{\mu, T}$ unless $\zeta^{\beta(b_1) - \mu_i} = \zeta^{\beta(b_2) - \mu_j}$, and then the action is well-defined unless also

$$\mu_i - \mu_j = d_{\beta(b_1)} - d_{\beta(b_2)} + r(\text{ct}(b_1) - \text{ct}(b_2)) c_0.$$

But then $\beta(b_1) = \beta(b_2)$ and $\text{ct}(b_1) = \text{ct}(b_2)$ implies $0 = \mu_i - \mu_j = S(b_1) - S(b_2)$, contradicting the column-strictness of S .

We write $e = \sum_{w \in S_n} w$ and

$$(2.5) \quad e = \sum_{w \in S_n} \sigma_w f_w$$

for some f_w in the ring obtained from $\mathfrak{t}$ by inverting $z_i - z_j$ for $1 \leq i < j \leq n$. Our first goal is to compute each f_w explicitly. Observe that $f_{w_0} = 1$ since no σ_w other than σ_{w_0} contributes w_0 . Next, for each $1 \leq i \leq n-1$ we have $s_i e = e$ and hence

$$\begin{aligned} \sum_{w \in S_n} \sigma_w f_w &= (\sigma_i - f_{\alpha_i}) \sum_{w \in S_n} \sigma_w f_w = \sum_{s_i w > w} \sigma_{s_i w} f_w + \sum_{s_i w < w} \sigma_i^2 \sigma_{s_i w} f_w - \sum_{w \in S_n} \sigma_w f_{w^{-1} \alpha_i} f_w \\ &= \sum_{s_i w < w} \sigma_w f_{s_i w} + \sum_{s_i w > w} \sigma_w f'_w - \sum_{w \in S_n} \sigma_w f_{w^{-1} \alpha_i} f_w \end{aligned}$$

for some $f'_w \in \mathfrak{t}$, where in the first and second summations we have reindexed, replacing w by $s_i w$. Thus for $w \in S_n$ with $s_i w < w$ comparing coefficients of σ_w gives the recurrence

$$(2.6) \quad f_w = f_{s_i w} - f_{w^{-1} \alpha_i} f_w \implies f_{s_i w} = (1 - f_{w^{-1} \alpha_i}) f_w.$$

Together with $f_{w_0} = 1$ this gives

$$(2.7) \quad f_w = \prod_{\substack{\alpha \in R^+ \\ \alpha \notin R(w)}} (1 - f_\alpha),$$

where

$$R(w) = \{\alpha \in R^+ \mid w(\alpha) \in R^-\}$$

is the inversion set of w . Hence

$$(2.8) \quad \sum_{w \in S_n} w = \sum_{w \in S_n} \sigma_w \prod_{\alpha \notin R(w)} (1 - f_\alpha).$$

Now, recalling that $g_\alpha \in F$ is defined by $f_\alpha \cdot f_{\mu,T} = g_\alpha f_{\mu,T}$, using (2.2) and the fact that σ_i is self-adjoint for $\langle \cdot, \cdot \rangle$ gives

$$(2.9) \quad \langle \sigma_i f_{\mu,T}, \sigma_i f_{\mu,T} \rangle = \langle f_{\mu,T}, \sigma_i^2 f_{\mu,T} \rangle = (1 - g_{\alpha_i}^2) \langle f_{\mu,T}, f_{\mu,T} \rangle$$

and hence

$$(2.10) \quad \langle \sigma_w f_{\mu,T}, \sigma_w f_{\mu,T} \rangle = \prod_{\alpha \in R(w)} (1 - g_\alpha^2) \langle f_{\mu,T}, f_{\mu,T} \rangle,$$

so that

$$\begin{aligned} \langle g_S, g_S \rangle &= \sum_{w \in S_n} \left(\prod_{\alpha \notin R(w)} (1 - g_\alpha) \right)^2 \langle \sigma_w(f_{\mu,T}), \sigma_w(f_{\mu,T}) \rangle \\ &= \sum_{w \in S_n} \left(\prod_{\alpha \notin R(w)} (1 - g_\alpha) \right)^2 \prod_{\alpha \in R(w)} (1 - g_\alpha^2) \langle f_{\mu,T}, f_{\mu,T} \rangle \\ &= n! \prod_{\alpha \in R^+} (1 - g_\alpha) \langle f_{\mu,T}, f_{\mu,T} \rangle \end{aligned}$$

where the last equality depends upon a specialization of the identity

$$\sum_{w \in S_n} \prod_{\epsilon_i - \epsilon_j \in R(w)} \left(1 + \frac{rc_0}{z_i - z_j} \right) \prod_{\epsilon_i - \epsilon_j \in R^+ - R(w)} \left(1 - \frac{rc_0}{z_i - z_j} \right) = n!$$

This last identity follows by (1) observing that the left-hand side is invariant by each simple reflection s_i , (2) observing that multiplying it by the discriminant clears its denominator, and (3) using the fact that the discriminant is the minimal degree alternating polynomial. In the formula for $\langle f_{\mu,T}, f_{\mu,T} \rangle$ from part (c) of Theorem 1.3 we reindex by setting $b = T^{-1}w_\mu(i)$ so that $\mu_i = S(b)$, and take products over $b \in \lambda^\bullet$ instead of $1 \leq i \leq n$. Combined with the preceding calculation, a bit of rearranging finishes the computation of $\langle g_S, g_S \rangle$. $\square$

3. THE MINIMAL DEGREE SYMMETRIC POLYNOMIAL AND ASPHERICAL VALUES

Throughout this section we fix an r -partition $\lambda^\bullet$ of n , and write S for the column-strict tableau on $\lambda^\bullet$ with $S(b) = l + (i-1)r$ if b is in the i th row of λ^l . The function g_S is the (unique up to scalars) minimal degree element of $M(\lambda^\bullet)^W$, and we set

$$(3.1) \quad g_{\lambda^\bullet} = g_S$$

with S as above. We have $g_S = e \cdot f_{\mu,T}$ for a Young tableau T that is uniquely determined by S , and where μ is the non-decreasing rearrangement of the entries of S . As in the proof of Theorem 2.2, we can construct $g_S = e \cdot f_{\mu,T}$ by rewriting e as a certain F -linear combination of intertwining operators σ_w , and it follows that the leading term of g_S is $n_\mu x^{\mu^+} w_0 v_T$, where n_μ is the size of the stabilizer

of μ in S_n . On the other hand, since the space of W -invariants in this degree is 1-dimensional by Theorem 2.2, $g_{\lambda^\bullet}$ must be a multiple of its specialization to $c_0 = d_0 = \cdots = d_{r-1} = 0$ (which is well-defined since the intertwiners σ_i have no poles when $c_0 = 0$). Since they have the same leading term they must be equal. Thus while any non-symmetric $f_{\mu,T}$ with $e.f_{\mu,T} = g_{\lambda^\bullet}$ may depend on the parameters, its symmetrization does not.

3.1. The norm of the minimal degree symmetric function. Let lim^l be the *lower rim* of λ^l , consisting of those boxes $b \in \lambda^l$ that are not directly above another box of λ^l . Similarly, let rim^l be the *right rim* of λ^l , consisting of those boxes which are not directly to the left of another box of λ^l . The *lower rim* $\text{lim}(\lambda^\bullet)$ and *right rim* $\text{rim}(\lambda^\bullet)$ of $\lambda^\bullet$ are the unions of the respective rims of its components. The *hook length product* associated to $\lambda^\bullet$ is

$$(3.2) \quad H_{\lambda^\bullet} = \prod_{\substack{b \in \text{lim}(\lambda^\bullet) \\ b' \in \text{rim}(\lambda^\bullet)}} \prod_{\substack{1 \leq k \leq S(b) - S(b') \\ k = \beta(b) - \beta(b') \pmod{r}}} (k - (d_{\beta(b)} - d_{\beta(b')}) - r(\text{ct}(b) - \text{ct}(b') - 1)c_0).$$

In the case $r = 1$ this reduces (after setting $c_0 = -\kappa$) to the hook length product from [Dun]. For each $0 \leq l \leq r - 1$ let b_{LL}^l be the lower left-hand corner box of λ^l , with the convention that if $\lambda^l = \emptyset$ then b_{LL}^l lies in the 0th row and 1st column, and put $S_l = S(b_{LL}^l)$ (so if $\lambda^l = \emptyset$ then $S_l = l - r$) and $c_l = \text{ct}(b_{LL}^l)$. Define an extra product $E_{\lambda^\bullet}$ by

$$(3.3) \quad E_{\lambda^\bullet} = \prod_{\substack{b \in \lambda^\bullet \\ 0 \leq l \leq r-1}} \prod_{\substack{1 \leq k \leq S(b) - S_l - r \\ k = \beta(b) - l \pmod{r}}} (k - (d_{\beta(b)} - d_l) - r(\text{ct}(b) - c_l + 1)c_0)$$

Theorem 3.1. *The norm of $g_{\lambda^\bullet}$ is given by*

$$\langle g_{\lambda^\bullet}, g_{\lambda^\bullet} \rangle = n! H_{\lambda^\bullet} E_{\lambda^\bullet}.$$

Proof. As in [Dun] we induct on the number of boxes in $\lambda^\bullet$. The base case with one box follows from Theorem 2.2. We sketch the inductive step, which is similar to that in [Dun]. In general, suppose $\lambda^\bullet$ is obtained from $\chi^\bullet$ by adding a box b with the property that $S(b) \geq S(b')$ for all $b' \in \chi^\bullet$. Then b is in particular a removable box, and Theorem 2.2 shows that

$$(3.4) \quad \begin{aligned} \langle g_{\lambda^\bullet}, g_{\lambda^\bullet} \rangle &= n \langle g_{\chi^\bullet}, g_{\chi^\bullet} \rangle \prod_{1 \leq k \leq S(b)} (k - (d_{\beta(b)} - d_{\beta(b)-k}) - r \text{ct}(b) c_0) \\ &\times \prod_{b' \in \chi^\bullet} \prod_{\substack{1 \leq k \leq S(b) - S(b') \\ k = \beta(b) - \beta(b') \pmod{r}}} \frac{k - (d_{\beta(b)} - d_{\beta(b')}) - r(\text{ct}(b) - \text{ct}(b') - 1)c_0}{k - (d_{\beta(b)} - d_{\beta(b')}) - r(\text{ct}(b) - \text{ct}(b'))c_0} \\ &\times \prod_{b' \in \chi^\bullet} \prod_{\substack{1 \leq k \leq S(b) - S(b') - r \\ k = \beta(b) - \beta(b') \pmod{r}}} \frac{k - (d_{\beta(b)} - d_{\beta(b')}) - r(\text{ct}(b) - \text{ct}(b') + 1)c_0}{k - (d_{\beta(b)} - d_{\beta(b')}) - r(\text{ct}(b) - \text{ct}(b'))c_0}. \end{aligned}$$

Fix a positive integer k and define $0 \leq l \leq r - 1$ by $k = \beta(b) - l \pmod{r}$. For each positive integer i with $k \leq S(b) - (l + (i - 1)r)$, the i th row of χ^l contributes a telescoping product totaling

$$(3.5) \quad \frac{k - (d_{\beta(b)} - d_l) - r(\text{ct}(b) - \text{ct}(b') - 1)c_0}{k - (d_{\beta(b)} - d_l) - r(\text{ct}(b) + i - 1)c_0}$$

to the second line in (3.4), where b' is the rightmost box in the i th row. Likewise, for each positive integer i with $k \leq S(b) - (l + (i - 1)r) - r$, the i th row of χ^l contributes a telescoping product totaling

$$(3.6) \quad \frac{k - (d_{\beta(b)} - d_l) - r(\text{ct}(b) + i)c_0}{k - (d_{\beta(b)} - d_l) - r(\text{ct}(b) - \text{ct}(b'))c_0}$$

to the third line in (3.4) where again b' is the rightmost box in the row. For $1 \leq i \leq \frac{1}{r}(S(b) - l - k)$ the denominator of (3.5) for the $i + 1$ th row and the numerator of (3.6) for the i th row cancel, and the denominator for the first row cancels with the factor $k - (d_{\beta(b)} - d_l) - \text{rct}(b)c_0$ from the first line of (3.4). The denominator of (3.6) cancels a corresponding factor in $H_{\lambda^\bullet}$ for the box directly above b in $\chi^\bullet$ (if there is one; otherwise the factor (3.6) does not appear), and the numerator of (3.6) for the bottom row of χ^l contributes to the expression for $E_{\lambda^\bullet}$ (if this bottom row consists of boxes b' with $S(b') < S(b)$, which can be the case only if $l \neq \beta(b) \bmod r$). The remaining factors in $E_{\lambda^\bullet}$ are those of the form

$$k - (d_{\beta(b)} - d_{\beta(b)-k}) - \text{rct}(b)c_0 \quad \text{for } k \text{ with } 1 \leq k \leq S(b) \text{ and } \lambda^{\beta(b)-k} = \emptyset.$$

These are accounted for by those factors in the first line of (3.4) that did not cancel with a denominator from (3.5). These observations together with the inductive hypothesis prove the formula. $\square$

3.2. Another formula for the norm. Here we record alternative expressions for $H_{\lambda^\bullet}$ and $E_{\lambda^\bullet}$ that are closer in spirit to the formulas of [DuOp] and [Dun]. Let ${}^t\mu$ denote the conjugate partition of the partition μ , and write

$$(x)_n = x(x+1)(x+2) \cdots (x+n-1)$$

for the Pochhammer symbol. If a partition $\mu = \emptyset$ let ${}^t\mu = (0)$, that is, ${}^t\mu_1 = 0$. In the following theorem, empty products are understood to equal 1.

Theorem 3.2. *There are $a_1, a_2 \in \mathbb{C}^\times$ with*

$$\begin{aligned} H_{\lambda^\bullet} = & a_1 \prod_{k=0}^{r-1} \prod_{l=0}^{k-1} \prod_{i=1}^{\min({}^t\lambda_1^k, {}^t\lambda_1^l)} \prod_{j=1}^{\lambda_i^k} \left(\frac{k-l}{r} - \frac{d_k-d_l}{r} + c_0 \left({}^t\lambda_j^k + \lambda_i^l - i - j + 1 \right) \right)_{t\lambda_j^k - i + 1} \\ & \times \prod_{l=k}^{r-1} \prod_{i=1}^{\min({}^t\lambda_1^k, {}^t\lambda_1^l)} \prod_{j=1}^{\lambda_i^k} \left(\frac{r+k-l}{r} - \frac{d_k-d_l}{r} + c_0 \left({}^t\lambda_j^k + \lambda_i^l - i - j + 1 \right) \right)_{t\lambda_j^k - i}. \end{aligned}$$

and

$$\begin{aligned} E_{\lambda^\bullet} = & a_2 \prod_{k=0}^{r-1} \prod_{l=0}^{k-1} \prod_{i=t\lambda_1^l+1}^{t\lambda_1^k} \prod_{j=1}^{\lambda_i^k} \left(\frac{k-l}{r} - \frac{d_k-d_l}{r} + c_0 \left(i - j - t\lambda_1^l \right) \right)_{i-t\lambda_1^l} \\ & \times \prod_{l=k}^{r-1} \prod_{i=t\lambda_1^l+2}^{t\lambda_1^k} \prod_{j=1}^{\lambda_i^k} \left(\frac{r+k-l}{r} - \frac{d_k-d_l}{r} + c_0 \left(i - j - t\lambda_1^l \right) \right)_{i-t\lambda_1^l-1}. \end{aligned}$$

Proof. The innermost product loops of our formulae can be expressed as Pochhammer symbols. For fixed k, l such that $1 \leq k, l \leq r-1$ the typical product has the form

$$P_{k,l}(U, A) = \prod_{\substack{1 \leq m \leq U \\ m \equiv k-l \pmod r}} (m - d_k + d_l - rAc_0).$$

Change the loop variable, letting $m = k - l + ri$, then i has integer values and satisfies $\frac{1-k+l}{r} \leq i \leq \frac{U-k+l}{r}$. Thus the upper limit is $u = \left\lfloor \frac{U-k+l}{r} \right\rfloor$, where we write $[x]$ for the largest integer at most x . If $0 \leq l \leq k-1$ then $-1 < -1 + \frac{2}{r} \leq \frac{1-k}{r} \leq \frac{1-k+l}{r} \leq 0$ and the lower limit is 0, and $P_{k,l}$ has $u+1$ terms; the vacuous product occurs for $u \leq -1$. If $k \leq l \leq r-1$ then $\frac{1}{r} \leq \frac{1+l-k}{r} \leq \frac{r-k}{r} \leq 1$ and the

lower limit is 1, and P has u terms (vacuous product for $u \leq 0$, but it is convenient to allow $u = 0$ in the following formula). In the latter case let $i = i' - 1$ then $0 \leq i' \leq u$; . Thus

$$P_{k,l}(U, A) = r^{u+1} \left(\frac{k-l}{r} - \frac{d_k - d_l}{r} - c_0 A \right)_{u+1}, 0 \leq l < k;$$

$$P_{k,l}(U, A) = r^u \left(\frac{r+k-l}{r} - \frac{d_k - d_l}{r} - c_0 A \right)_u, k \leq l \leq r-1.$$

Consider the product $H_{\lambda^\bullet}$. The typical $b \in \text{Irim}(\lambda^\bullet)$ has $\beta(b) = k$, coordinates $({}^t\lambda_j^k, j)$ for $1 \leq j \leq \lambda_1^k$ and $S(b) = k + r({}^t\lambda_j^k - 1)$, $\text{ct}(b) = j - {}^t\lambda_j^k$. The typical $b' \in \text{rrim}(\lambda^\bullet)$ has $\beta(b) = l$, coordinates (i, λ_i^l) for $1 \leq i \leq {}^t\lambda_1^l$ and $S(b') = l + r(i - 1)$, $\text{ct}(b') = \lambda_i^l - i$. The corresponding term in $H_{\lambda^\bullet}$ is $P_{k,l}(k - l + r({}^t\lambda_j^k - i), j - {}^t\lambda_j^k + i - \lambda_i^l - 1)$. Evaluate $u = \lfloor \frac{U-k+l}{r} \rfloor = {}^t\lambda_j^k - i$. The condition ${}^t\lambda_j^k - i \geq 0$ is equivalent to $\lambda_i^l - j \geq 0$. This imposes the additional bound $i \leq {}^t\lambda_1^k$, and thus the pertinent index values are $1 \leq i \leq \min({}^t\lambda_1^k, {}^t\lambda_1^l)$, $1 \leq j \leq {}^t\lambda_i^k$. These arguments deduce the formula for $H_{\lambda^\bullet}$ from the one in the Theorem.

Consider the product $E_{\lambda^\bullet}$. The typical $b \in \lambda^\bullet$ has $\beta(b) = k$, coordinates (i, j) with $1 \leq i \leq {}^t\lambda_1^k$, $1 \leq j \leq \lambda_i^k$ and $S(b) = k + r(i - 1)$, $\text{ct}(b) = j - i$. The corresponding b_{LL} has $\beta(b_{LL}) = l$, $S(b_{LL}) = l + r({}^t\lambda_1^l - 1)$, $\text{ct}(b_{LL}) = 1 - {}^t\lambda_1^l$, so that $U = k - l + r(i - {}^t\lambda_1^l) - 1$ and $u = i - {}^t\lambda_1^l - 1$ and $A = j - i + {}^t\lambda_1^l$. Each pair (i, j) with ${}^t\lambda_1^l + 1 \leq i \leq {}^t\lambda_1^k$, $1 \leq j \leq \lambda_i^k$ contributes to $E_{\lambda^\bullet}$ (when $k \leq l$ and $i = {}^t\lambda_1^l + 1$ the contribution to the product is 1). This demonstrates the alternative formula for $E_{\lambda^\bullet}$. $\square$

3.3. Aspherical values. In this subsection we will, for the first time, specialize parameters to complex numbers.

A parameter $c = (c_0, d_0, \dots, d_{r-1})$ is *aspherical* if $\mathbb{H}_c e_+ \mathbb{H}_c \neq \mathbb{H}_c$, where $e_+ = \frac{1}{|W|} \sum_{w \in W} w$ is the trivial idempotent for $W = G(r, 1, n)$. By Theorem 4.1 of [BeEt], c is aspherical exactly if there is an r -partition $\lambda^\bullet$ of n so that $L_c(\lambda^\bullet)^W = 0$.

We thank Pavel Etingof for pointing out the following lemma and that it follows from S. Montarani's work [Mon]. We give a proof based on [Gri] together with [BeEt].

Lemma 3.3. (a) *If there is a rectangle λ with at most n boxes, an integer $0 \leq l \leq r-1$, and a positive integer k with*

$$k = d_l - d_{l-k} + r \text{ct}(b) c_0, \quad k \not\equiv 0 \pmod{r}, \quad \text{and} \quad 1 \leq k \leq l + (\text{row}(b) - 1)r$$

where b is the removable (lower right hand corner) box of λ , then c is aspherical.

(b) *Parameters c such that $c_0 = -k/m$ for integers $1 \leq k < m \leq n$ are aspherical.*

Proof. (a) Consider the module $M_c(\lambda^\bullet)$ for the rational Cherednik algebra of type $G(r, 1, m)$, where m is the number of boxes in the rectangle and $\lambda^\bullet$ is the r -partition with $\lambda^l = \lambda$ and all other components empty. Provided that c_0 is not a rational number, Theorem 7.5 of [Gri] implies that the span of the $f_{\mu, T}$ for $T \in \text{SYT}(\lambda^\bullet)$ and $\mu_i \geq k$ for some $1 \leq i \leq m$ is a submodule of $M_c(\lambda^\bullet)$; in particular, by the inequality $k \leq l + (\text{row}(b) - 1)r$ all the polynomials g_S are in this submodule. Therefore $\langle g, h \rangle = 0$ for all symmetric functions $g, h \in M_c(\lambda^\bullet)^W$. By continuity this condition continues to hold for all parameters c on the hyperplane, implying that such c are aspherical for the rational Cherednik algebra for $G(r, 1, m)$. By Theorem 4.1 of [BeEt], it is aspherical also for the rational Cherednik algebra of type $G(r, 1, n)$.

(b) These points are (some of the) zeros of the norm of $g_{\lambda^\bullet}$, where $\lambda^{r-1} = (1, 1, \dots, 1)$ and $\lambda^l = \emptyset$ for $l \neq r-1$ corresponds to the determinant representation of $G(r, 1, n)$. So they are aspherical because every invariant in $M(\lambda^\bullet)$ is a multiple of $g_{\lambda^\bullet}$. $\square$

In other words, in addition to the hyperplanes $c_0 = -k/m$ with $1 \leq k < m \leq n$, for each integer m with $-(n-1) \leq m \leq n-1$ the hyperplanes

$$(3.7) \quad k = d_l - d_{l-k} + rmc_0 \quad \text{with} \quad 1 \leq k \leq l + \left(\sqrt{n + \frac{1}{4}m^2} - \frac{1}{2}m - 1 \right) r$$

are also aspherical.

Theorem 3.4. *The set of aspherical values c for the rational Cherednik algebra for $G(r, 1, n)$ is the union of the hyperplanes $c_0 = -k/m$ for integers $1 \leq k < m \leq n$ and hyperplanes*

$$k = d_l - d_{l-k} + rct(b)c_0$$

where b is the lower right-hand corner box of a rectangle with at most n boxes, $0 \leq l \leq r-1$, $k \not\equiv 0 \pmod{r}$, and $1 \leq k \leq l + (\text{row}(b) - 1)r$.

Proof. By Lemma 3.3 the stated hyperplanes consist of aspherical values. Conversely, assume that $c = (c_0, d_0, \dots, d_{r-1})$ is aspherical and choose an r -partition $\lambda^\bullet$ of n so that $L_c(\lambda^\bullet)^W = 0$.

Writing $g_{\lambda^\bullet} = \sum g_T v_T$ for certain $g_T \in \mathbb{C}[x_1, \dots, x_n]$ we have

$$\langle g_{\lambda^\bullet}, g_{\lambda^\bullet} \rangle = \sum \langle v_T, \overline{g_T} \cdot g_{\lambda^\bullet} \rangle,$$

where $\overline{\cdot}$ is the W -equivariant conjugate linear isomorphism of $\mathbb{C}[x_1, \dots, x_n]$ onto $\mathbb{C}[y_1, \dots, y_n]$ determined by $\overline{x_i} = y_i$. While the left hand side is only defined for real values of the parameters (as otherwise the contravariant form does not specialize appropriately), the right hand side always makes sense and is a polynomial in the parameters $(c_0, d_0, \dots, d_{r-1})$. For generic real values of the parameters it is given by the formula in Theorem 3.1. It follows that it is given by this formula for all values of the parameters.

Thus four situations can occur, corresponding to the zeros of the formula in Theorem 3.1:

- (a) If one of the factors of $H_{\lambda^\bullet}$ corresponding to boxes b, b' with $\beta(b) = \beta(b')$ is zero, then number c_0 is rational, of the form $c_0 = -k/m$ with integers $1 \leq k < m \leq n$.
- (b) If one of the factors in $E_{\lambda^\bullet}$ corresponding to an empty λ^j gives a zero, then there is a box $b \in \lambda^\bullet$ and an integer $1 \leq k \leq S(b)$ (necessarily $\beta(b) = S(b) \not\equiv j \pmod{r}$) with $k = \beta(b) - j \pmod{r}$ such that

$$k = d_{\beta(b)} - d_{\beta(b)-k} + rct(b)c_0,$$

in which case putting $l = \beta(b)$ and taking the rectangle with corner at b does the trick.

- (c) There are boxes $b \in \text{lim}(\lambda^\bullet)$ and $b' \in \text{rim}(\lambda^\bullet)$ and an integer k with $1 \leq k \leq S(b) - S(b')$ and $0 \not\equiv k = \beta(b) - \beta(b') \pmod{r}$ so that

$$k = d_{\beta(b)} - d_{\beta(b')} - r(ct(b) - ct(b') - 1)c_0.$$

Let $l = \beta(b)$ and let $m = ct(b) - ct(b') - 1$. Using (3.7) it will suffice to show that

$$S(b) - S(b') \leq l + \left(\sqrt{n + \frac{m^2}{4}} - \frac{m}{2} - 1 \right) r.$$

Write

$$x = \text{row}(b), \quad y = \text{col}(b), \quad x' = \text{row}(b'), \quad \text{and} \quad y' = \text{col}(b').$$

Since $S(b) - S(b') = l - \beta(b') + (x - x')r$ it suffices to show that

$$x - x' \leq \sqrt{n + \frac{(y - x - y' + x' - 1)^2}{4}} - \frac{y - x - y' + x' - 1}{2} - 1,$$

or, after rearranging, that

$$(x - x' + 1)(y - y') \leq n.$$

This inequality follows from $xy \leq n$ and $x'y' \leq n$.

- (d) There is a box $b \in \lambda^\bullet$, an integer $0 \leq i \leq r-1$ so that $\lambda^i \neq \emptyset$ and if b' is the lower left hand corner of λ^i then $S(b') < S(b)$, and an integer $1 \leq k \leq S(b) - S(b') - r$ with $0 \neq k = \beta(b) - \beta(b') \bmod r$ and

$$k = d_{\beta(b)} - d_{\beta(b')} + r(\text{ct}(b) - \text{ct}(b') + 1)c_0.$$

Setting $m = \text{ct}(b) - \text{ct}(b') + 1$ it suffices to show that

$$S(b) - S(b') - r \leq l + \left(\sqrt{n + \frac{m^2}{4}} - \frac{m}{2} - 1 \right) r,$$

and the rest of the proof of this case proceeds as in case (c). □

3.4. Other linear characters. Let W be a complex reflection group, and with the notation of section 5 of Rouquier's paper [Rou] let $\mathbb{H}_h$ be the rational Cherednik algebra with parameter h attached to W . Fix a linear character $\xi : W \rightarrow \mathbb{C}^\times$ and write $e_\xi = \sum_{w \in W} \xi(w)^{-1} w$ for the corresponding symmetrizer. With θ_ξ as in section 3.3.1 of [Rou], there is an isomorphism $\mathbb{H}_h \rightarrow \mathbb{H}_{\theta_\xi(h)}$ such that $e_\xi \mapsto e_{\text{triv}}$, and it follows that $\mathbb{H}_h e_\xi \mathbb{H}_h \neq \mathbb{H}_h$ exactly if $\mathbb{H}_{\theta_\xi(h)} e_{\text{triv}} \mathbb{H}_{\theta_\xi(h)} \neq \mathbb{H}_{\theta_\xi(h)}$.

In case $W = G(r, 1, n)$, a character ξ is determined by the integers $0 \leq i \leq 1$ and $0 \leq j \leq r-1$ with $\xi(s_1) = (-1)^i$ and $\xi(\zeta_1) = \zeta^j$. In terms of our parametrization, the parameter $\theta_\xi(h)$ is obtained by replacing c_0 by $(-1)^i c_0$ and d_l by d_{l+j} . We deduce the following corollary.

Corollary 3.5. *Let ξ be the linear character of $G(r, 1, n)$ determined by $\xi(s_1) = (-1)^i$ and $\xi(\zeta_1) = \zeta^j$. A parameter c is such that $\mathbb{H}_c e_\xi \mathbb{H}_c \neq \mathbb{H}_c$ exactly if one of the following holds:*

- (a) $c_0 = (-1)^{i+1} k/m$ for integers $1 \leq k < m \leq n$, or
- (b) there are integers $0 \leq l \leq r-1$ and k and a rectangle with at most n boxes so that writing b for its lower right-hand corner box,

$$1 \leq k \leq l + (\text{row}(b) - 1)r \quad \text{and} \quad k = d_{l+j} - d_{l+j-k} + (-1)^i r \text{ct}(b) c_0.$$

3.5. The groups $G(r, p, n)$. Fix a positive integer p dividing r and let $G(r, p, n)$ be the subgroup of $G(r, 1, n)$ consisting of matrices so that the product of the non-zero entries is an r/p -th root of unity. When $n \geq 3$ (to avoid fusion of conjugacy classes of reflections) and $d_i = d_j$ for $i = j \bmod r/p$, the rational Cherednik algebra for $G(r, p, n)$ is the subalgebra of that for $G(r, 1, n)$ generated by $\mathbb{C}[x_1, \dots, x_n]$, $G(r, p, n)$, and $\mathbb{C}[y_1, \dots, y_n]$. As in section 9 of [Gri], it is the fixed subalgebra for a cyclic group of automorphisms of $\mathbb{H}_c$, which allows one to apply (an especially simple version of) Clifford theory to relate representations of the two algebras.

Let C be the cyclic shift (by r/p) operator on r -partitions given by

$$C.(\lambda^0, \dots, \lambda^{r-1}) = (\lambda^{r/p}, \lambda^{r/p+1}, \dots, \lambda^{r/p-1}).$$

Thus the representations of $G(r, 1, n)$ whose restriction to $G(r, p, n)$ contains the trivial representation are indexed by precisely those r -partitions in the C -orbit of $((n), \emptyset, \dots, \emptyset)$.

With notation as in section 9 of [Gri], the equation

$$L_c(\lambda^\bullet) = \bigoplus_{q=0}^{p/k-1} L_c(\lambda^\bullet, q)$$

implies that $L_c(\lambda^\bullet, q)^{G(r, p, n)} = 0$ for some q exactly if $L_c(\lambda^\bullet)^\xi = 0$ for all linear characters ξ of $G(r, 1, n)$ that restrict to the trivial linear character of $G(r, p, n)$.

With the specialization $d_i = d_j$ when $i = j \bmod r/p$, Corollary 3.5 implies that c is ξ -aspherical for some ξ restricting trivially to $G(r, p, n)$ exactly if it is aspherical. Therefore c is aspherical for the rational Cherednik algebra of type $G(r, p, n)$ exactly if it is so for $G(r, 1, n)$. We obtain:

Corollary 3.6. *For $n \geq 3$, the aspherical set for the rational Cherednik algebra of type $G(r, p, n)$ is given by the equations of Theorem 3.4 restricted to parameters with $d_i = d_j$ for $i = j \bmod r/p$.*

4. ORDERING CATEGORY $\mathcal{O}_c$

4.1. The ordering, numerically. For a non-zero number $m \in \mathbb{R}^\times$ and $a, b \in \mathbb{R}$ write $a = b \bmod m$ if $(a - b)m^{-1} \in \mathbb{Z}$. For multisets X and Y of real numbers write $X = Y \bmod m$ if the corresponding multisets of equivalence classes are equal. Let $c = (c_0, d_1, \dots, d_{r-1}) \in \mathbb{R}^r$ be a parameter with $c_0 > 0$ and define a partial order $\geq_c$ on the set of r -partitions of n by the rule: $\lambda^\bullet \geq_c \chi^\bullet$ if for all $j \in \mathbb{R}$ and $0 \leq l \leq r - 1$,

$$(4.1) \quad \begin{aligned} & |\{b \in \lambda^\bullet \mid \frac{d_{\beta(b)}}{rc_0} + \text{ct}(b) > j \text{ or } \frac{d_{\beta(b)}}{rc_0} + \text{ct}(b) = j \text{ and } \beta(b) \leq l\}| \\ & \geq |\{b' \in \chi^\bullet \mid \frac{d_{\beta(b')}}{rc_0} + \text{ct}(b') > j \text{ or } \frac{d_{\beta(b')}}{rc_0} + \text{ct}(b') = j \text{ and } \beta(b') \leq l\}|. \end{aligned}$$

Also define an equivalence relation $\equiv_c$ by $\lambda^\bullet \equiv_c \chi^\bullet$ if there is an equality of multisets

$$(4.2) \quad \{\text{ct}(b) + \frac{d_{\beta(b)} - \beta(b)}{rc_0} \mid b \in \lambda^\bullet\} = \{\text{ct}(b) + \frac{d_{\beta(b)} - \beta(b)}{rc_0} \mid b \in \chi^\bullet\} \bmod c_0^{-1}.$$

For the definition of a highest weight category, see [Rou], Definition 4.11. The congruence condition part of the following theorem is analogous to a result of [GrLe] concerning cyclotomic Hecke algebras. In [LyMa] Lyle and Mathas finished the classification of the blocks of cyclotomic Hecke algebras (and showed that they are the same as those for the affine Hecke algebra). It might be that the norm formulas presented in this paper are enough to give an alternative proof of this theorem; we have not yet attempted to do so.

Theorem 4.1. *For a parameter $c = (c_0, d_1, \dots, d_{r-1}) \in \mathbb{R}^r$ with $c_0 > 0$ category $\mathcal{O}_c$ is a highest weight category with respect to the order given by $\lambda^\bullet \geq \chi^\bullet$ if $\lambda^\bullet \geq_c \chi^\bullet$ and $\lambda^\bullet \equiv_c \chi^\bullet$.*

Proof. We will show that if $[M_c(\lambda^\bullet) : L_c(\chi^\bullet)] \neq 0$ then $\lambda^\bullet \geq_c \chi^\bullet$ and $\lambda^\bullet \equiv_c \chi^\bullet$. Assuming this for the moment, for each r -partition $\lambda^\bullet$, write $P_c(\lambda^\bullet)$ for the projective cover of $L_c(\lambda^\bullet)$. By Corollary 2.10 of [GGOR] $M_c(\lambda^\bullet)$ is a quotient of $P_c(\lambda^\bullet)$. It follows from the formula following Theorem 2.19 and Proposition 3.3 of [GGOR] that if $[P_c(\chi^\bullet) : M_c(\lambda^\bullet)] \neq 0$ then $\lambda^\bullet \geq \chi^\bullet$, and hence $\mathcal{O}_c$ is a highest weight category with respect to the order $\geq$.

If $[M_c(\lambda^\bullet) : L_c(\chi^\bullet)] \neq 0$ then upon examining generalized $\mathfrak{t}$ -eigenspaces part (a) of Theorem 1.3 implies that there exist orderings $b_1, b_2, \dots, b_n$ and $b'_1, b'_2, \dots, b'_n$ of the boxes of $\lambda^\bullet$ and $\chi^\bullet$, respectively, and non-negative integers μ_i so that

$$(4.3) \quad \beta(b_i) - \mu_i = \beta(b'_i) \bmod r$$

and

$$(4.4) \quad \mu_i = d_{\beta(b_i)} - d_{\beta(b'_i)} + r(\text{ct}(b_i) - \text{ct}(b'_i))c_0$$

for $1 \leq i \leq n$. The latter condition can be rewritten

$$(4.5) \quad 0 \leq \mu_i = \left(\frac{d_{\beta(b_i)}}{rc_0} + \text{ct}(b_i) - \left(\frac{d_{\beta(b'_i)}}{rc_0} + \text{ct}(b'_i) \right) \right) rc_0,$$

and setting $\mu_i = q_i r + \beta(b_i) - \beta(b'_i)$ one obtains also

$$(4.6) \quad 0 \leq q_i = \left(\frac{d_{\beta(b_i)} - \beta(b_i)}{rc_0} + \text{ct}(b_i) - \left(\frac{d_{\beta(b'_i)} - \beta(b'_i)}{rc_0} + \text{ct}(b'_i) \right) \right) c_0$$

This implies $\lambda^\bullet \equiv_c \chi^\bullet$, and also since $c_0 > 0$, equations (4.5) and (4.6) imply that for $1 \leq i \leq n$

$$(4.7) \quad \frac{d_{\beta(b_i)}}{rc_0} + \text{ct}(b_i) \geq \frac{d_{\beta(b'_i)}}{rc_0} + \text{ct}(b'_i) \quad \text{with equality implying } \beta(b_i) \leq \beta(b'_i).$$

Hence for all $j \in \mathbb{R}$ and $0 \leq l \leq r-1$,

$$\begin{aligned} & |\{b \in \lambda^\bullet \mid \frac{d_{\beta(b)}}{rc_0} + \text{ct}(b) > j \text{ or } \frac{d_{\beta(b)}}{rc_0} + \text{ct}(b) = j \text{ and } \beta(b) \leq l\}| \\ & \geq |\{b' \in \chi^\bullet \mid \frac{d_{\beta(b')}}{rc_0} + \text{ct}(b') > j \text{ or } \frac{d_{\beta(b')}}{rc_0} + \text{ct}(b') = j \text{ and } \beta(b') \leq l\}|, \end{aligned}$$

which is (4.1). $\square$

4.2. Cores, quotients, and beta numbers. The basic reference we use for this material is Chapter 1 of [Mac]. See especially exercise 8 of section 1.

Let s be a complex number and let λ be a partition. The set of *beta numbers* of λ with respect to s is

$$(4.8) \quad B_s(\lambda) = \{\lambda_j + s - j + 1 \mid 1 \leq j < \infty\}.$$

We are using the notation B_s to avoid a conflict with the function β we defined on boxes of an r -partition. The first $\ell(\lambda)$ beta numbers of λ are $s+1$ more than the contents of the rightmost boxes of λ . Since no two of the rightmost boxes of λ can have the same content, λ and s can be reconstructed from the beta numbers.

The reflection representation of $S_r = W(A_{r-1})$ is

$$\{(a_1, a_2, \dots, a_r) \in \mathbb{R}^r \mid \sum_{i=1}^r a_i = 0\} \text{ and the root lattice is } Q = \{(a_1, \dots, a_r) \in \mathbb{Z}^r \mid \sum_{i=1}^r a_i = 0\}.$$

Let $\lambda^\bullet = (\lambda^0, \lambda^1, \dots, \lambda^{r-1})$ be an r -partition and let $a = (a_1, a_2, \dots, a_r)$ be an element of the root lattice Q of type A_{r-1} . As in Section 6 of [Gor], we will construct from this data a partition λ whose r -quotient is $\lambda^\bullet$ and whose r -core is determined by a . In order to conform with the notation in Gordon's paper we set $\lambda^{(i)} = \lambda^{r-i}$, for $1 \leq i \leq r$. The set of B_0 numbers of λ is

$$(4.9) \quad B_0(\lambda) = \bigcup_{1 \leq i \leq r} \{i + r(x-1) \mid x \in B_{a_i}(\lambda^{(i)})\}.$$

This defines a bijection

$$(4.10) \quad Q \times \Pi^r \rightarrow \Pi,$$

where Π is the set of partitions.

4.3. Dominance order. If λ and χ are (ordinary) partitions of n then we write $\lambda \geq \chi$ if for all $1 \leq i \leq n$ we have

$$(4.11) \quad \sum_{j=1}^i \lambda_j \geq \sum_{j=1}^i \chi_j.$$

The relation $\geq$ is called *dominance order*. Note that we denoted it by $\geq_d$ in Section 1 in order to distinguish it from the partial order we defined on $\mathbb{Z}_{\geq 0}$, but since the latter ordering plays no role from now on we henceforth drop the subscript d and reserve the symbol $\geq$ for dominance order.

The following lemma gives the relationship between dominance order and the special case $r=1$ of Theorem 4.1; we will use it to prove Corollary 4.3.

Lemma 4.2. *Suppose λ and χ are partitions of n . Then $\lambda \geq \chi$ if and only if there exist numbering $b_1, b_2, \dots, b_n$ and $b'_1, b'_2, \dots, b'_n$ of the boxes of λ and χ such that $\text{ct}(b_i) \geq \text{ct}(b'_i)$ for $1 \leq i \leq n$. Equivalently, $\lambda \geq \chi$ if and only if for each $j \in \mathbb{Z}$, one has*

$$(4.12) \quad |\{b \in \lambda \mid \text{ct}(b) \geq j\}| \geq |\{b' \in \chi \mid \text{ct}(b') \geq j\}|$$

Proof. Suppose $\lambda \geq \chi$. By [Mac] (1.15) we may obtain λ from χ by applying a sequence of raising operators (that is, operators that move a single box to a higher row). But when $\lambda = R\chi$ for a single raising operator R the equation (4.12) is clear.

Conversely, assuming that numberings as in the statement of the lemma exist, it follows that for each $c \in \mathbb{Z}$ there are at least as many boxes of λ with content at least c as boxes of χ with content at least c . For $1 \leq i \leq n$ such that $\lambda_i > 0$, let $c_i = \lambda_i - i$ be the content of the last box in the i th row of λ and compute

$$\begin{aligned} \sum_{j=1}^i (\lambda_j - \chi_j) &= |\{\text{boxes in first } i \text{ rows of } \lambda \text{ with content at least } c_i\}| \\ &\quad - |\{\text{boxes in first } i \text{ rows of } \chi \text{ with content at least } c_i\}| \\ &\quad + |\{\text{boxes in first } i \text{ rows of } \lambda \text{ with content less than } c_i\}| \\ &\quad - |\{\text{boxes in first } i \text{ rows of } \chi \text{ with content less than } c_i\}| \\ &\geq |\{\text{boxes in } \lambda \text{ with content at least } c_i\}| \\ &\quad - |\{\text{boxes in } \chi \text{ with content at least } c_i\}| \geq 0. \end{aligned}$$

Here we used the fact that

$$|\{\text{boxes in first } i \text{ rows of } \lambda \text{ with content at least } c_i\}| = |\{\text{boxes in } \lambda \text{ with content at least } c_i\}|.$$

□

If λ is a partition then for $k \in \mathbb{Z}$ and $s \in \mathbb{R}$, the relationship between beta numbers and contents described in 4.2 implies

$$(4.13) \quad |\{b \in \lambda \mid \text{ct}(b) = k\}| = \begin{cases} |\{x \in B_s(\lambda) \mid x \geq k + s + 1\}| & \text{if } k \geq 0, \text{ and} \\ |\{x \in B_s(\lambda) \mid x \geq k + s + 1\}| + k & \text{if } k < 0. \end{cases}$$

We set $a_i = d_{r-i}/rc_0$ and assume for the remainder of this section that $a_i \in \mathbb{Z}$ for $1 \leq i \leq r$. If λ is the partition corresponding to $\lambda^\bullet$ and $a = (a_1, \dots, a_r)$ under the bijection of Section 4.2, then for $j \in \mathbb{R}$ using (4.13) gives

$$\begin{aligned} |\{b \in \lambda \mid \text{ct}(b) \geq j\}| &= \sum_{\substack{k \in \mathbb{Z} \\ k \geq j}} \sum_{\substack{1 \leq l \leq r}} |\{x \in B_{a_l}(\lambda^{(l)}) \mid x \geq (k - l + 1)/r + 1\}| + \sum_{k \in \mathbb{Z}, j \leq k < 0} k \\ &= \sum_{\substack{k \in \mathbb{Z} \\ k \geq j}} \sum_{\substack{1 \leq l \leq m_k + 1}} |\{x \in B_{a_l}(\lambda^{(l)}) \mid x \geq (k - m_k)/r + 1\}| \\ &\quad + \sum_{m_k + 1 < l \leq r} |\{x \in B_{a_l}(\lambda^{(l)}) \mid x \geq (k - m_k)/r\}| + \sum_{k \in \mathbb{Z}, j \leq k < 0} k \end{aligned}$$

where for $k \in \mathbb{Z}$ we write m_k for the integer with $0 \leq m_k \leq r-1$ and $k = m_k \bmod r$. Therefore using (4.13) again

$$\begin{aligned}
|\{b \in \lambda \mid \text{ct}(b) \geq j\}| &= \sum_{\substack{k \in \mathbb{Z} \\ k \geq j}} \sum_{1 \leq l \leq m_k+1} |\{b \in \lambda^{(l)} \mid \text{ct}(b) = (k - m_k)/r - a_l\}| \\
&+ \sum_{m_k+1 < l \leq r} |\{b \in \lambda^{(l)} \mid \text{ct}(b) = (k - m_k)/r - a_l - 1\}| + \sum_{k \in \mathbb{Z}, j \leq k < 0} k \\
&- \sum_{\substack{k \in \mathbb{Z}, k \geq j \\ 1 \leq l \leq m_k+1 \\ (k-m_k)/r - a_l < 0}} ((k - m_k)/r - a_l) - \sum_{\substack{k \in \mathbb{Z}, k \geq j \\ m_k+1 < l \leq r \\ (k-m_k)/r - a_l - 1 < 0}} ((k - m_k)/r - a_l - 1)
\end{aligned}$$

Note that the last three summands depend on the sequence $a_1, \dots, a_r$ and the number j but are independent of λ . Write $f(a_\bullet, j)$ for their sum and $n_j = rq_j + m_j$ with n_j the least integer greater than or equal to j and q_j, m_j integers with $0 \leq m_j \leq r-1$. In the sums over k above we reindex by setting $q = (k - m_k)/r$ to obtain

$$\begin{aligned}
|\{b \in \lambda \mid \text{ct}(b) \geq j\}| - f(a_\bullet, j) &= \sum_{m=m_j+1}^r \sum_{q \geq q_j} \left(\sum_{l=1}^m |\{b \in \lambda^{(l)} \mid \text{ct}(b) = q - a_l\}| + \sum_{l=m+1}^r |\{b \in \lambda^{(l)} \mid \text{ct}(b) = q - a_l - 1\}| \right) \\
&+ \sum_{m=1}^{m_j} \sum_{q \geq q_j+1} \left(\sum_{l=1}^m |\{b \in \lambda^{(l)} \mid \text{ct}(b) = q - a_l\}| + \sum_{l=m+1}^r |\{b \in \lambda^{(l)} \mid \text{ct}(b) = q - a_l - 1\}| \right) \\
&= \sum_{m=1}^r \sum_{q \geq q_j+1} \left(\sum_{l=1}^m |\{b \in \lambda^{(l)} \mid \text{ct}(b) = q - a_l\}| + \sum_{l=m+1}^r |\{b \in \lambda^{(l)} \mid \text{ct}(b) = q - a_l - 1\}| \right) \\
&+ \sum_{m=m_j+1}^r \left(\sum_{l=1}^m |\{b \in \lambda^{(l)} \mid \text{ct}(b) = q_j - a_l\}| + \sum_{l=m+1}^r |\{b \in \lambda^{(l)} \mid \text{ct}(b) = q_j - a_l - 1\}| \right).
\end{aligned}$$

Counting the number of appearances of each summand of the form $|\{b \in \lambda^{(l)} \mid \text{ct}(b) = X\}|$ for each integer $X \geq q_j - 1 - a_l$ gives

$$\begin{aligned}
|\{b \in \lambda \mid \text{ct}(b) \geq j\}| - f(a_\bullet, j) &= r \sum_{l=1}^r |\{b \in \lambda^{(l)} \mid \text{ct}(b) \geq q_j + 1 - a_l\}| + \sum_{l=1}^r (l-1) |\{b \in \lambda^{(l)} \mid \text{ct}(b) = q_j - a_l\}| \\
&+ \sum_{1 \leq l \leq m_j+1} (r - m_j) |\{b \in \lambda^{(l)} \mid \text{ct}(b) = q_j - a_l\}| + \sum_{m_j+1 < l \leq r} (r - l + 1) |\{b \in \lambda^{(l)} \mid \text{ct}(b) = q_j - a_l\}| \\
&+ \sum_{m_j+1 < l \leq r} (l - m_j - 1) |\{b \in \lambda^{(l)} \mid \text{ct}(b) = q_j - 1 - a_l\}| \\
&= r \sum_{l=1}^r |\{b \in \lambda^{(l)} \mid \text{ct}(b) + a_l \geq q_j + 1\}| + \sum_{1 \leq l \leq m_j+1} (r - m_j - 1 + l) |\{b \in \lambda^{(l)} \mid \text{ct}(b) + a_l = q_j\}| \\
&+ r \sum_{m_j+1 < l \leq r} |\{b \in \lambda^{(l)} \mid \text{ct}(b) + a_l = q_j\}| + \sum_{m_j+1 < l \leq r} (l - m_j - 1) |\{b \in \lambda^{(l)} \mid \text{ct}(b) + a_l = q_j - 1\}|
\end{aligned}$$

which, recalling $\lambda^l = \lambda^{(r-l)}$ and $a_l = d_{r-l}/rc_0$ and reindexing by interchanging l and $r-l$, is equal to

$$\begin{aligned}
&= r \sum_{l=0}^{r-1} |\{b \in \lambda^l \mid \text{ct}(b) + d_l/rc_0 \geq q_j + 1\}| + \sum_{r-m_j-1 \leq l \leq r-1} (2r - m_j - 1 - l) |\{b \in \lambda^l \mid \text{ct}(b) + d_l/rc_0 = q_j\}| \\
&+ r \sum_{0 \leq l < r-m_j-1} |\{b \in \lambda^l \mid \text{ct}(b) + d_l/rc_0 = q_j\}| \\
&+ \sum_{0 \leq l < r-m_j-1} (r - m_j - 1 - l) |\{b \in \lambda^l \mid \text{ct}(b) + d_l/rc_0 = q_j - 1\}| \\
&= \sum_{0 \leq l < r-m_j-1} |\{b \in \lambda^\bullet \mid \text{ct}(b) + d_{\beta(b)}/rc_0 > q_j - 1, \text{ or } = q_j - 1 \text{ and } \beta(b) \leq l\}| \\
&+ \sum_{r-m_j-1 \leq l \leq r-1} |\{b \in \lambda^\bullet \mid \text{ct}(b) + d_{\beta(b)}/rc_0 > q_j, \text{ or } = q_j \text{ and } \beta(b) \leq l\}|.
\end{aligned}$$

The final equality above is obtained by counting the number of times a given box $b \in \lambda^\bullet$ contributes, according to the cases $\text{ct}(b) + d_{\beta(b)}/rc_0 = q_j - 1$, $\text{ct}(b) + d_{\beta(b)}/rc_0 = q_j$ and $\beta(b) < r - m_j - 1$ or $\beta(b) \geq r - m_j - 1$, and $\text{ct}(b) + d_{\beta(b)}/rc_0 \geq q_j + 1$.

Write $\lambda^\bullet \geq'_c \chi^\bullet$ if $\lambda \geq \chi$ in dominance order; note that λ and χ depend on c via the bijection described above. The preceding calculation and Theorem 4.1 give the answer to Question 10.1 of Gordon's paper [Gor]:

Corollary 4.3. *Assume that $c_0 > 0$ and $d_l/rc_0 \in \mathbb{Z}$ for $0 \leq l \leq r - 1$. Category $\mathcal{O}_c$ for $\mathbb{H}_c$ is a highest weight category with respect to the order $\geq'_c$.*

REFERENCES

- [BeCh] Y. Berest and O. Chalykh, *Quasi-invariants of complex reflection groups*, arXiv:0912.4518v1
- [BeEt] R. Bezrukavnikov and P. Etingof, *Parabolic induction and restriction functors for rational Cherednik algebras*, Selecta Math. (N.S.) 14 (2009), no. 3-4, 397–425. arXiv:0803.3639
- [BEG] Y. Berest, P. Etingof, and V. Ginzburg, *Cherednik algebras and differential operators on quasi-invariants*, Duke Math. J. 118 (2003), no. 2, 279–337. arXiv:math/0111005
- [Dun] C. Dunkl, *Symmetric and antisymmetric vector-valued Jack polynomials*, arXiv:1001.4485
- [DuOp] C. Dunkl and E. Opdam, *Dunkl operators for complex reflection groups*, Proc. London Math. Soc. (3) 86 (2003), no. 1, 70–108. arXiv:math/0108185
- [GGS] V. Ginzburg, I. Gordon, and T. Stafford, *Differential operators and Cherednik algebras*, Selecta Math. (N.S.) 14 (2009), no. 3-4, 629–666.
- [GGOR] Ginzburg, Guay, Opdam, and Rouquier, *On the category $\mathcal{O}$ for rational Cherednik algebras*, Invent. Math. 154 (2003), no. 3, 617–651. arXiv:math/0212036
- [Gor] I. Gordon, *Quiver varieties, category $\mathcal{O}$ for rational Cherednik algebras, and Hecke algebras*, Int. Math. Res. Pap. IMRP 2008, no. 3, Art. ID rpn006, 69 pp. arXiv:math/0703150
- [GoSt] I. Gordon and J.T. Stafford, *Rational Cherednik algebras and Hilbert schemes*, Adv. Math. 198 (2005), no. 1, 222–274, arXiv:math/0407516
- [GrLe] J. Graham and G. Lehrer, *Cellular algebras* Invent. Math. 123 (1996), no. 1, 1–34.
- [Gri] S. Griffeth, *Orthogonal functions generalizing Jack polynomials*, to appear in Transactions of the American Mathematical Society, arXiv:0707.0251
- [Gri2] S. Griffeth, *Towards a combinatorial representation theory for the rational Cherednik algebra of type $G(r, p, n)$* , to appear in Proceedings of the Edinburgh Mathematical Society, arXiv:math/0612733v3
- [LyMa] S. Lyle and A. Mathas, *Blocks of cyclotomic Hecke algebras*, Adv. Math. 216 (2007), no. 2, 854–878, arXiv:math/0607451
- [Mac] I. Macdonald, *Symmetric functions and Hall polynomials*, second edition, Oxford Mathematical Monographs, 1995.
- [Mac2] I. Macdonald, *Affine Hecke algebras and orthogonal polynomials*, Cambridge University Press, 2003.
- [Mon] S. Montarani, *On some finite dimensional representations of symplectic reflection algebras associated to wreath products*, Comm. Algebra 35 (2007), no. 5, 1449–1467. arXiv:math/0411286

- [Rou] R. Rouquier, *q-Schur algebras and complex reflection groups*, Mosc. Math. J. 8 (2008), no. 1, 119–158, 184.
arXiv:math/0509252

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, P.O. Box 400137, CHARLOTTESVILLE, VA 22904-4137,

E-mail address: `cf5z@virginia.edu`

SCHOOL OF MATHEMATICS, UNIVERSITY OF EDINBURGH, EDINBURGH EH9 3JZ

E-mail address: `griffeth@ed.ac.uk`